Linker Nucleoside, Its Preparation and Use INVENTOR(S): Miculka et al. U.S. Serial No.: Not yet assigned Page: 1 of 4 Exp Mail label EV337189665US


1/4

Fig. 1.

Page: 2 of 4 Exp Mail label EV337189665US

2/4

Fig. 2A



$$B = A^{Bz} = \frac{1}{N} \frac{HHBz}{H}$$

$$B^{Bz} = A^{(Bz)} = \frac{1}{N} \frac{H}{N}$$

$$B^{Bz} = A^{(Bz)} = \frac{1}{N} \frac{H}{N}$$

$$B^{Bz} = U^{Bz} = \frac{1}{N} \frac{1}{N} \frac{1}{N} \frac{1}{N}$$

$$B^{Bz} = U^{Bz} = \frac{1}{N} \frac{1}{N} \frac{1}{N} \frac{1}{N} \frac{1}{N}$$

Bz = C (Dimethylamino) Pyridine

CH3)2HPy = 4-(Dimethylamino) Pyridine

DMI = 4.4'-Dimethoxytrityl

BSA = Bis(trimethylsityl) acetamide

CPG = 'long-chain-alkylamino-controlled

pore glass' (Sigma)

Fig. 2B

Linker Nucleoside, Its Preparation and Use INVENTOR(S): Miculka et al.
U.S. Serial No.: Not yet assigned Page: 4 of 4
Exp Mail label EV337189665US

4/4




Fig. 3